

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Ciprofloxacin Tablets USP safely and effectively. See full prescribing information for Ciprofloxacin Tablets, USP (ciprofloxacin hydrochloride) tablet, for oral use Initial U.S. Approval: 1987

WARNING: SERIOUS ADVERSE REACTIONS INCLUDING TENDINITIS, TENDON RUPTURE, PERIPHERAL NEUROPATHY, CENTRAL NERVOUS SYSTEM EFFECTS AND EXACERBATION OF MYASTHENIA GRAVIS

- Fluoroquinolones, including Ciprofloxacin, have been associated with disabling and potentially irreversible serious adverse reactions that have occurred together (5.1), including:
 - Tendinitis and tendon rupture (5.2)
 - Peripheral neuropathy (5.3)
 - Central nervous system effects (5.4)

Discontinue Ciprofloxacin immediately and avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who experience any of these serious adverse reactions (5.1)

- Fluoroquinolones, including Ciprofloxacin, may exacerbate muscle weakness in patients with myasthenia gravis. Avoid Ciprofloxacin in patients with known history of myasthenia gravis (5.5)
- Because fluoroquinolones, including Ciprofloxacin, have been associated with serious adverse reactions (5.1-5.15), reserve Ciprofloxacin for use in patients who have no alternative treatment options for the following indications:
 - Acute exacerbation of chronic bronchitis (1.10)
 - Acute uncomplicated cystitis (1.11)
 - Acute sinusitis (1.12)

RECENT MAJOR CHANGES

Warnings and Precautions (5.15) 7/2017

INDICATIONS AND USAGE

Ciprofloxacin Tablets 250 mg, 500 mg, and 750 mg are a fluoroquinolone antibacterial indicated in adults (≥18 years of age) with the following infections caused by designated, susceptible bacteria and in pediatric patients where indicated:

- Skin and Skin Structure Infections (1.1)
- Bone and Joint Infections (1.2)
- Complicated Intra-Abdominal Infections (1.3)
- Infectious Diarrhea (1.4)
- Typhoid Fever (Enteric Fever) (1.5)
- Uncomplicated Cervical and Urethral Gonorrhea (1.6)
- Inhalational Anthrax post-exposure in adult and pediatric patients (1.7)
- Plague in adult and pediatric patients (1.8)
- Chronic Bacterial Prostatitis (1.9)
- Lower Respiratory Tract Infections (1.10)
- Acute Exacerbation of Chronic Bronchitis (1.11)
- Urinary Tract Infections (1.11)
- Urinary Tract Infections (UTI) or Cystitis (1.12)
- Complicated UTI and Pyelonephritis in Pediatric Patients (1.12)
- Acute Sinusitis (1.12)

Usage To reduce the development of drug-resistant bacteria and maintain the effectiveness of Ciprofloxacin and other antibacterial drugs, Ciprofloxacin should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria (1.13)

DOSE AND ADMINISTRATION

Adult Dosage Guidelines			
Infection	Dose	Frequency	Duration
Skin and Skin Structure	500-750 mg	every 12 hours	7 to 14 days
Bone and Joint	500-750 mg	every 12 hours	4 to 8 weeks
Complicated Intra-Abdominal	500 mg	every 12 hours	7 to 14 days
Infectious Diarrhea	500 mg	every 12 hours	5 to 7 days
Typhoid Fever	500 mg	every 12 hours	10 days
Uncomplicated Gonorrhea	250 mg	single dose	single dose
Inhalational anthrax (post-exposure)	500 mg	every 12 hours	60 days
Plague	500-750 mg	every 12 hours	14 days
Chronic Bacterial Prostatitis	500 mg	every 12 hours	28 days

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: SERIOUS ADVERSE REACTIONS INCLUDING TENDINITIS, TENDON RUPTURE, PERIPHERAL NEUROPATHY, CENTRAL NERVOUS SYSTEM EFFECTS AND EXACERBATION OF MYASTHENIA GRAVIS

- INDICATIONS AND USAGE
- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Bone and Joint Infections
- Complicated Intra-Abdominal Infections
- Infectious Diarrhea
- Typhoid Fever (Enteric Fever)
- Uncomplicated Cervical and Urethral Gonorrhea
- Inhalational Anthrax (Post-Exposure)
- Plague
- Chronic Bacterial Prostatitis
- Lower Respiratory Tract Infections
- Urinary Tract Infections
- Acute Sinusitis

- Usage
- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

- Disabling and Potentially Irreversible Serious Adverse Reactions Including Tendinitis and Tendon Rupture, Peripheral Neuropathy, and Central Nervous System Effects and Exacerbation of Myasthenia Gravis
- Urinary Tract Infections
- Acute Sinusitis

Adult Dosage Guidelines			
Infection	Dose	Frequency	Duration
Lower Respiratory Tract	500-700 mg	every 12 hours	7 to 14 days
Urinary Tract	250-500 mg	every 12 hours	7 to 14 days
Acute Uncomplicated Cystitis	250 mg	every 12 hours	3 days
Acute Sinusitis	500 mg	every 12 hours	10 days

- Adults with creatinine clearance 30-50 mL/min 250-500 mg q 12 h (2.3)
- Adults with creatinine clearance ≤29 mL/min 250-500 mg q 18 h (2.3)
- Patients on hemodialysis or peritoneal dialysis 250-500 mg q 24 h (after dialysis) (2.3)

Pediatric Oral Dosage Guidelines			
Infection	Dose	Frequency	Duration
Complicated UTI and Pyelonephritis (1 to 17 years of age)	10-20 mg/kg (maximum 750 mg per dose)	Every 12 hours	10-21 days
Inhalational Anthrax (Post-Exposure)	15 mg/kg (maximum 500 mg per dose)	Every 12 hours	60 days
Plague	15 mg/kg (maximum 500 mg per dose)	Every 8 to 12 hours	10-21 days

DOSE FORMS AND STRENGTHS

- Tablets: 250 mg, 500 mg, 750 mg (3)

CONTRAINDICATIONS

- Known hypersensitivity to Ciprofloxacin or other quinolones (4.1, 5.6, 5.7)
- Concomitant administration with tizanidine (4.2)

WARNINGS AND PRECAUTIONS

- Hypersensitivity and other serious reactions: Serious and sometimes fatal reactions (for example, anaphylactic reactions) may occur after the first or subsequent doses of Ciprofloxacin. Discontinue Ciprofloxacin at the first sign of skin rash, jaundice or any sign of hypersensitivity. (4.1, 5.6, 5.7)
- Hepatotoxicity: Discontinue immediately if signs and symptoms of hepatitis occur. (5.8)
- Clostridium difficile*-associated diarrhea: Evaluate if colitis occurs. (5.10)
- QT Prolongation: Prolongation of the QT interval and isolated cases of torsade de pointes have been reported. Avoid use in patients with known prolongation, those with hypokalemia, and with other drugs that prolong the QT interval. (5.11, 7.7, 8.5)
- Inhalational Anthrax post-exposure in adult and pediatric patients (1.7)
- Plague in adult and pediatric patients (1.8)
- Chronic Bacterial Prostatitis (1.9)
- Lower Respiratory Tract Infections (1.10)
- Acute Exacerbation of Chronic Bronchitis (1.11)
- Urinary Tract Infections (1.11)
- Urinary Tract Infections (UTI) or Cystitis (1.12)
- Complicated UTI and Pyelonephritis in Pediatric Patients (1.12)
- Acute Sinusitis (1.12)

The most common adverse reactions ≥ 1% were nausea, diarrhea, liver function tests abnormal, vomiting, and rash. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Carlsbad Technology, Inc. at 1-855-397-9777 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Interacting Drug	Interaction
Theophylline	Serious and fatal reactions. Avoid concomitant use. Monitor serum level (7)
Warfarin	Anticoagulant effect enhanced. Monitor prothrombin time, INR, and bleeding (7)
Antidiabetic agents	Hypoglycemia including fatal outcomes have been reported. Monitor blood glucose (7)
Phenyltin	Monitor phenyltin level (7)
Methotrexate	Monitor for methotrexate toxicity (7)
Cyclosporine	May increase serum creatinine. Monitor serum creatinine (7)

Decreased Ciprofloxacin absorption. Take 2 hours before or 6 hours after Ciprofloxacin (7)

As with other drugs, some isolates of *Pseudomonas aeruginosa* may develop resistance fairly rapidly during treatment with ciprofloxacin. Culture and susceptibility testing performed periodically during therapy will provide information not only on the therapeutic effect of the antimicrobial agent but also on the possible emergence of bacterial resistance.

Ciprofloxacin Tablets should be administered orally as described in the appropriate Dosage Guidelines table.

The determination of dosage and duration for any particular patient must take into consideration the severity and nature of the infection, the susceptibility of the causative microorganism, the integrity of the patient's host-defense mechanisms, and the status of renal and hepatic function.

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide. Revised: 03/2018

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

Because fluoroquinolones, including Ciprofloxacin, have been associated with serious adverse reactions (see Warnings and Precautions (5.1-5.15)) and for some patients ADRs are limiting, Ciprofloxacin for treatment of AECB in patients who have no alternative treatment options.

1.11 Urinary Tract Infections

Ciprofloxacin is indicated in adult patients for treatment of urinary tract infections caused by *Escherichia coli*, *Klebsiella pneumoniae*, *Enterobacter cloacae*, *Serratia marcescens*, *Proteus mirabilis*, *Providencia rettgeri*, *Morganella morganii*, *Citrobacter koseri*, *Citrobacter freundii*, *Pseudomonas aeruginosa*, methicillin-susceptible *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, or *Enterococcus faecalis*.

Ciprofloxacin is indicated in adult female patients for treatment of acute uncomplicated cystitis caused by *Escherichia coli* or *Staphylococcus saprophyticus*.

Because fluoroquinolones, including Ciprofloxacin, have been associated with serious adverse reactions (see Warnings and Precautions (5.1-5.15)) and for some patients acute uncomplicated cystitis is self-limiting, reserve Ciprofloxacin for treatment of acute uncomplicated cystitis in patients who have no alternative treatment options.

Ciprofloxacin is indicated in pediatric patients aged one to 17 years of age for treatment of complicated urinary tract infections (cUTI) and pyelonephritis due to *Escherichia coli* (see Use in Specific Populations (8.4)).

Although effective in clinical trials, Ciprofloxacin is not a drug of first choice in the pediatric population due to an increased incidence of adverse reactions compared to controls, including reactions related to joints and/or surrounding tissues. Ciprofloxacin, like other fluoroquinolones, is associated with arthropathy and histopathological changes in the joints of juvenile animals (see Warnings and Precautions (5.12), Adverse Reactions (6.1), Use in Specific Populations (8.4) and Nonclinical Toxicology (13.2)).

Ciprofloxacin is indicated in adult patients for treatment of acute sinusitis caused by *Haemophilus influenzae*, *Streptococcus pneumoniae*, or *Moraxella catarrhalis*.

Because fluoroquinolones, including Ciprofloxacin, have been associated with serious adverse reactions (see Warnings and Precautions (5.1-5.15)) and for some patients acute sinusitis is self-limiting, reserve Ciprofloxacin for treatment of acute sinusitis in patients who have no alternative treatment options.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Ciprofloxacin and other antibacterial drugs, Ciprofloxacin should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

If anaerobic organisms are suspected of contributing to the infection, appropriate therapy should be initiated. Appropriate culture and susceptibility tests should be performed before treatment in order to isolate and identify organisms causing infection and to determine their susceptibility to Ciprofloxacin. Therapy with Ciprofloxacin may be initiated before results of these tests are known; once results become available appropriate therapy should be continued.

As with other drugs, some isolates of *Pseudomonas aeruginosa* may develop resistance fairly rapidly during treatment with ciprofloxacin. Culture and susceptibility testing performed periodically during therapy will provide information not only on the therapeutic effect of the antimicrobial agent but also on the possible emergence of bacterial resistance.

Ciprofloxacin Tablets should be administered orally as described in the appropriate Dosage Guidelines table.

The determination of dosage and duration for any particular patient must take into consideration the severity and nature of the infection, the susceptibility of the causative microorganism, the integrity of the patient's host-defense mechanisms, and the status of renal and hepatic function.

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

See full prescribing information for use in pediatric and geriatric patients (8.4, 8.5)

Pediatric patients with moderate to severe renal insufficiency were excluded from the clinical trial of cUTI and pyelonephritis. No information is available on the extent to which Ciprofloxacin for treatment of AECB in patients who have renal insufficiency (that is, creatinine clearance < 50 mL/min/1.73m²).

2.4 Important Administration Instructions

Concomitant administration of Ciprofloxacin with dairy products (like milk or yogurt) or calcium-fortified juices alone should be avoided since decreased absorption is possible; however, Ciprofloxacin may be taken with a meal that contains these products.

Hydration of Patients Receiving Ciprofloxacin Assure adequate hydration of patients receiving Ciprofloxacin to prevent the formation of highly concentrated urine. Crystalluria has been reported with quinolones.

Instruct the patient of the appropriate Ciprofloxacin administration (see Patient Counseling Information (17)).

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

Discontinue Ciprofloxacin immediately at the first signs or symptoms of any serious adverse reaction. In addition, avoid the use of fluoroquinolones, including Ciprofloxacin, in patients who have experienced any of these serious adverse reactions associated with fluoroquinolones.

